

异叶苣苔属 (苦苣苔科) 雌蕊的畸形发育及其系统意义*

王印政** 潘开玉 李振宇 洪德元

(中国科学院植物研究所系统与进化植物学开放研究实验室, 北京 100093)

AN ANAMORPHOSIS OF GYNOECEIUM IN *WHYTOCKIA* (GESNERIACEAE), WITH PHYLOGENETIC IMPLICATION*

WANG Yin-Zheng**, PAN Kai-Yu, LI Zhen-Yu, HONG De-Yuan

(Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences,
Beijing 100093)

关键词 畸形发育, 返祖现象, 雌蕊, 异叶苣苔属

Key words Anamorphosis, Atavism, Gynoecium, *Whytockia*

The genus *Whytockia* W. W. Smith, mainly characterized by anisophylly, bilocular ovary with axile placentation and ebracteate pair-flowered cymes, is considered as the most primitive representative of the relict tribe Klugieae (Weber, 1982). Obviously, it is an ancient relic of the stock from which the other genera of Klugieae have evolved. But the origin and the ancestral group of its own is left unknown. An anamorphosis in the ontogeny of gynoecium found in *Whytockia* seems significant in explaining the origin of the gynoecium in *Whytockia*.

MATERIALS AND METHODS

Buds and flowers at different developmental stages of the two species—*Whytockia hekouensis* Y. Z. Wang and *W. bijensis* Y. Z. Wang in the study were obtained from Hekou of Yunnan and Bijie of Guizhou, and fixed in FAA. The material used for SEM was dehydrated, critical point dried and Au / Pd sputter coated. Serial sections were stained with safranin and fast green.

RESULTS AND DISCUSSION

As shown in Plate I: 4, the abnormal young gynoecium in *Whytockia hekouensis* consisted of two opening and unequal carpels which were born at different levels on the receptacle. Two whorls of the bracts surrounded the gynoecium, and also there existed some appendages between the gynoecium and bracts. The two carpels respectively folded adaxially and involutely along their median axis where the enlarged contacts

• Supported by a Special Grant from the President of Chinese Academy of Sciences

• • present address: Kunming Institute of Botany, CAS, Heilongtan, Kunming 650204, Yunnan

formed through the fusion of each two margins of carpels. The lower carpel was arch-shaped as a result of the unequal development while the upper carpel was like a hemi-tube opening upward. The epidermal hairs grew out from inside of the lower carpel. Contrasting sharply with the abnormal gynoeceium, the normal gynoeceium demonstrated an almost synchronous development, and an early closure and fusion of the two carpels, which had initiated from a pentagonal primordium through the developmental stage of two horse-shoe-shaped carpel flanges (Plate I: 1~3).

As indicated in the results, each carpel in the anamorphosis of the gynoeceium in *Whytockia* resembles the early development of a carpel in an apocarpous gynoeceium which is homologous with an involute leaf (Fahn, 1982). In addition, the relict characteristics in *Whytockia* are also related to the anamorphosis. The evidences are the carpellary margins being fused in the lower part but free in the middle part of the bilocular ovary (Plate I: 6,7), and the epidermal hairs often observed in the septal surfaces and inner wall of ovary (Plate I: 5). In general, the ventral edges may be viewed as corresponding to the contact region in an involute leaf (Eames, 1961). The carpel in the ancestor of *Whytockia* may have originated directly from a highly involute procarpel in which the contact is along the margins of procarpel in the lower part, but back on the abaxial surface of procarpel in the middle part where the carpellary margins may be free and inside the locule.

Thus, we can conclude that the anamorphosis of the gynoeceium in *Whytockia* is a reversible atavism in which some characteristics have lost in the modern plant of *Whytokia* such as the opening and involute carpels born at different levels on the receptacle, while some have maintained such as the epidermal hairs within the ovary and the free carpellary margins. The bicarpellate syncarpous gynoeceium in *Whytokia* may have evolved from the ancestor with two apocarpous gynoeceium born alternately on the receptacle and originated directly from a highly involute closure of the procarpel. The phylogenetic condensation of the gynoeceial internode, and later the fusion between the two apocarpous gynoeceia may have taken place in the origin of the genus *Whytockia*. Moreover, the bracts, appendages and the unequal carpels in the atavism are respectively parallel to the two whorls of perianth, the stamens and the anisophyllous leaves in the normal plant of *Whytockia*. Whether there exists a phylogenetic relation between them still needs further comparative studies in the fields of teratology, ontogeny and evolutionary morphology.

REFERENCES

- Eames A J, 1961. Morphology of the angiosperms. New York: McGraw-Hill, 186~245
 Fahn A, 1982. Plant Anatomy. Oxford: Pergamon Press, 386~426
 Weber A, 1982. Contributions to the morphology and systematics of Klugieae and Loxonieae (Gesneriaceae): IX.
 The genus *Whytockia*. Notes from the Royal Botanic Garden Edinburgh 40(2): 359~367

Explanation of Plates

1~3. The initiation and early development of the normal gynoeceium in *W. hekouensis*. 4. The abnormal young gynoeceium surrounded by two whorls of bracts and some appendages in *W. hekouensis*. 5. Part of the cross section of ovary in *W. bijiensis*. 6~7. The cross section in the lower and middle part of ovary in *W. hekouensis*.


C1, C2: primordia of the two carpels at different developmental stages. cr: contact region of an involute carpel. B1, B2: two whorls of bracts. col: column. →: indicating the epidermal hairs.

王印政等: 异叶苣苔属雌蕊的畸形发育及其系统意义

图版 I

WANG Yin-Zheng *et al.*: An Anamorphosis of Gynoccium in *Whytockia*
(Gesneriaceae), with Phylogenetic implication

Plate I



See explanation at the end of text